

Infraspinatus tendon release



One of the most useful and efficient ways to convey information to you about your pet is via the written word. We carefully craft these notes to give you helpful information and accurate expectations around your pet's surgical experience.

Please read this. *Please save it and read it during each stage of the recovery process.* Surprises make for a stressful time for everyone.

Photo and video examples may be found on our website: www.directvetsurg.com in the Pet Owner Portal under "DVS Resources".

Your pet has had shoulder surgery with removal of abnormal scarring (i.e. contracture) of the infraspinatus tendon. The defect in the tendon has been removed and the restriction of the shoulder joint has been released; the shoulder joint will return to normal range of motion and function. Patients generally return to normal leg performance after the healing period.

GENERAL INFORMATION

Please keep a note of your questions as you and your pet progress thru recovery and address them to your primary care veterinary team. ****Use your checklist at the end of these notes to make these check-ins complete.****

First few days postop

MONITORING

Please keep your pet in a comfortable, safe, indoor location without free access to stairs for the next 24 hours as he/she recovers from anesthesia and surgery.

Your pet may be groggy for the next few days. He or she may whine or appear more anxious than usual; this may indicate pain/discomfort or side-effects of the medications. Please call your primary care veterinary team for assistance with medication adjustments or return for exam and additional pain medications as needed.

Monitor appetite and attitude. *If both do not steadily improve over the next 2-3 days,* please call your primary care veterinary team or return for progress evaluation and problem-solving.

You can expect your pet to have a bowel movement within 5 days. Some animals take longer than others depending on when they last ate prior to surgery and when they started eating after surgery. It may be abnormal in color and consistency for 2-3 days. If you have any concerns, please speak with your primary care veterinary team.

Please confirm that your pet has urinated within 24 hours of returning home. If he/she does not, or you notice any problems related to urination, please speak with your primary care veterinary team.

Closely supervise your pet's movements over the first 3-4 days when s/he will be groggy from anesthetics and pain medications. Do not allow access to dangerous situations/locations (i.e. stairs, furniture, small children, other pets). Limit activities to necessary events only (on leash/out to go to the bathroom; access to food/water.)

Please look at the incision when you get home; knowing what "normal" looks like is helpful later when trying to understand if the incision is healing abnormally. Photos are helpful for you and us, when communicating from a distance.

MEDICATIONS

It is likely that you have been prescribed one or more medications (given by mouth) for your pet over the first 2-3wks of recovery. During the discharge appointment or shortly thereafter, please make sure you understand:

- what each medication is being used to treat,
- what side-effects may develop, and
- whether or not the medication should be refilled and continued.

BANDAGE CARE (IF PRESENT)

A clear, plastic bandaid may have been applied to the incision (Tegaderm). This will protect the incision from infection from the environment and a wayward pet tongue! It is useful for 2-3 days. You may remove it like a "bandaid" at any time.

Week 1-2 postop

MONITORING

Please look at the incision twice daily. It should be dry, slightly red along the margins, slightly swollen/thick on the edges with a light crust down the center. Over several days, it should lose redness and swelling.

Problems to call your veterinarian about:

- gapping (the edges should be exactly touching)
- ongoing or new discharge (other than small amount of crusting)
- swelling (other than slightly raised skin near edges).

Some bruising is normal and will resolve in 5-7 days.

The occasional patient will have more extensive bruising and/or a puffy ankle on day 2-3. If this develops, please let your primary care veterinarian know; a visit or sending photos will help them characterize the issue and decide on a course of action. The vast majority of the time, this is a transient issue that will follow a typical 5-7 day course resolving.

It is not uncommon for a seroma to develop under the incision; this is a pocket of normal tissue fluids that develops in a high motion area such as the shoulder joint. If the swelling

is smaller than a plum, please monitor. Warm compresses applied to the area will encourage the fluid to reabsorb more quickly (3-7 days.) If the seroma progressively enlarges, please have your primary care veterinarian assess the problem.

Do not allow your pet to lick or chew the incision. Pets tend to want to lick early in the healing period and scratch later in the healing period; this can compromise the incision and predispose to infection. If necessary, please prevent access to the incision by using creative clothing options (ex. long-sleeved T-shirt), an E-collar or other devices, if you must leave your pet unattended. See www.directvetsurg.com, Pet Owner portal→Pet Links and DVS Resources

TEST PENDING

If there were any samples removed for testing, they will be submitted and results usually will be available in 5-7 days. Your primary care team will contact you with test results and make any treatment changes as needed.

PROGRESS EXAMS

Please return to your primary care clinic in 10-14 days for a progress exam. Skin healing will be evaluated, sutures (if present) will be removed, use of the limb will be assessed and any questions you have will be addressed. (Alternatively, you can schedule a remote progress exam to include a photo of the incision and a videoclip of your pet walking around the room. Include a brief summary for the veterinary team and any questions you have.)

Your pet should start touching his/her toe down within the first 2 weeks. Thereafter, leg use should steadily improve to 90% normal at 6-8 weeks. If you notice a sudden deterioration in leg use at any time after surgery, please see your veterinarian for exam.

DIET

The **most** beneficial and **least** expensive treatment for joint health (and general health) over the lifetime of your pet is maintaining your pet on the lean/thin side of normal his/her whole life. Any orthopedic condition can progress with arthritis over time due to excessive wear & tear; carrying less body weight will relieve some of this stress from the joints. Good parameters to monitor body condition are:

- 1) you should be able to feel the ribs and pelvic bones, but not see them;
- 2) your pet should have an "hour glass" figure when viewed from above looking down;
- 3) your pet should have a tucked-up belly when viewed from the side.

RESTRICTIONS

Restrict your pet from any rigorous activity for 4 weeks. No running, jumping, playing. For dogs, short, leashed walks around the yard to urinate/defecate are fine. Walking around one level of the house is fine. Prevent all pets from jumping up and down from furniture, etc. A bad landing or an awkward take-off can result in serious repercussions.

Confine your pet, when unattended, to one level/section of the house on carpeted floors or crate confine. Rugs can be used to reduce the risk of slipping on slippery surfaces.

Limited, supervised access to stairs is recommended for 8 weeks. Restrict access to stairs when your pet is unattended (baby gates, etc.) Use caution and direct supervision on stairs (up and down) with a hand on the collar and a safety strap/leash under the belly to prevent falls.

Please **always** use a short (6 ft) leash when taking your pet outside to urinate/defecate during this restriction period. Use a belly band/sling/strap for safety when walking across slippery floors & icy sidewalks to prevent falling

If your pet is not using the limb by 2 weeks, if you notice a sudden deterioration, or if your pet stops using the leg at any time after surgery, please call your veterinarian for advice. While it is not unusual for brief episodes of discomfort over 2-3 months of healing, limping that continues for more than 24 hours should be examined. X-rays or other tests may be needed to diagnose significant problems.

The *Help 'em Up harness* (see company website; or similar) is a very useful tool that your dog can wear during the restricted period that allows you to have a quick "handle" to grasp by the shoulders to assist your pet to rise and lie down, climb/descend stairs, save from a slippery floor, etc.

If your pet's personality-type challenges these restrictions, discuss options for medicinal-assistance for your pet to make restrictions easier to manage (i.e. Trazadone, Gabapentin, Tramadol, other).

PHYSICAL THERAPY REGIMEN

Our lives are often very busy, so if you must err, err on the "do less" side of these instructions. Less physical therapy will result in a slower return to function, but more aggressive physical therapy by a non-professional too early may result in failure of the surgical repair. Cats (and some dogs) often resist physical therapy dramatically; avoid any activity that results in major uncooperative behavior.

Studies have shown that a formal PT program can decrease post-operative recovery time. Please let us or your veterinarian know if you are interested in a professional physical therapy referral in the Twin Cities.

Week 1: Apply ice packs (wrapped in thin cloth) to incision area twice daily for 10-15 minutes. Baggies of frozen peas work well for this, or make an ice pack by freezing 2 parts isopropyl alcohol to one part water in a ziplock bag. Continue 5 days.

Week 2: Range of Motion Exercise- Have your pet lie on his/her good side. Apply a warm compress to the shoulder, and hold with one hand. Hold the elbow with the other hand. Slowly and gently push the arm backward into full flexion of shoulder; hold for 5 seconds. Slowly pull the arm forward into full extension of the shoulder; hold for 5 seconds. Repeat

this motion 15-20 times slowly once to twice daily. This exercise should not be performed to the point of pain or resentment.

Weeks 3-6: Swimming is wonderful rehabilitation exercise when performed correctly. You may allow controlled swimming after week 2. Controlled swimming requires that your pet not jump or leap into the water; walking into the water until it is deep enough to swim is required. Throwing balls to fetch often results in sudden jumping and lunging, which can cause serious problems in the healing phase. Do not over extend your pet; start with short excursions (5 minutes) and increase duration and frequency gradually. Do not exceed once daily swimming until week 6.

Long-term lifestyle

The prognosis for dogs treated surgically for infraspinatus tendon contracture is considered good to excellent. The majority of dogs return to a normal gait, level of activity, and endurance. Following the 4 week restricted period, there are no recommended limitations to their lifestyle.

Checklist:

HOME MONITORING AND PROGRESS CHECK-IN WITH VETERINARY TEAM

- Pet attitude and appetite—Are these improving daily? What are your observations? What are your specific concerns?
- Pet mobility—Is this improving daily? What are your observations? What are your specific concerns?
- Incision health—Is redness and swelling going away? Is there discharge or moisture? (Photos taken close-up and at different angles are helpful for your primary care veterinary team,)
- Pet pain level—What are your observations? What are your specific concerns?
- Other

Based on your own experience through this, we welcome and encourage suggestions to this information that may help future patients and their people. Pay it forward! (directvetsurg@gmail.com)

--The DVS Crew